

- cerned with the evolution of a communal breeding system. *Folia Primatol.* 42: 17-32.
- Goldizen, A. W. and Terborgh, J. 1989. Demography and dispersal patterns of a tamarin population: possible causes of delayed breeding. *Am. Nat.* 134: 208-224.
- Hampton, J. K., Jr., Hampton, S. H. and Landwehr, B. T. 1966. Observations on a successful breeding colony of the marmoset, *Oedipomidas oedipus*. *Folia Primatol.* 4: 265-287.
- Hernández-Camacho, J. and Cooper, R. W. 1976. The nonhuman primates of Colombia. In: *Neotropical Primates: Field Studies and Conservation*, R. W. Thorington, Jr. and P. G. Heltne (eds.), pp.35-69. National Academy of Sciences, Washington, D. C.
- Jämmrich, S. 1985. Zum Fortpflanzungsverhalten des Weißbüscheläffchens, *Callithrix jacchus* (L., 1758). *Zool. Garten (N.F.)* 55: 177.
- Kleiman, D. G. 1977. Monogamy in mammals. *Quart. Rev. Biol.* 52: 39-69.
- Kleiman, D. G. 1986. The sociobiology of captive propagation. In: *Conservation Biology. The Science of Scarcity and Diversity*. Soulé, M. E. (ed.), pp. 243-261. Sinauer Associates, Inc., Sunderland, Massachusetts.
- König, A. and Siess, M. 1986. Zwei Fälle von vaterloser Zwillingsaufzucht bei *Callithrix penicillata* x *jacchus* und *Callithrix geoffroyi*. *Zool. Garten (N.F.)* 56: 438-440.
- Leutenegger, W. 1980. Monogamy in callitrichids: A consequence of phyletic dwarfism? *Int. J. Primatol.* 1 (1): 95-98.
- Price, E. C. and McGrew, W. C. 1991. Departures from monogamy in colonies of captive cotton-top tamarins. *Folia Primatol.* 57: 16-27.
- Rothe, H. 1978. Sub-grouping behaviour in captive *Callithrix jacchus* families: a preliminary investigation. In: *Biology and Behaviour of Marmosets*, H. Rothe, H.-J. Wolters and J. P. Hearn (eds.), pp.233-257. Eigenverlag H. Rothe, Göttingen.
- Rothe, H. and Darms, K. 1993. The social organization of marmosets: a critical evaluation of recent concepts. In: *Marmosets and Tamarins: Systematics, Behaviour, and Ecology*, A. B. Rylands (ed.), pp.1476-199. Oxford University Press, Oxford.
- Rothe, H., Darms, K., König, A., Siess, M. and Brusek, P. 1986. Dynamics of group size in the common marmoset (*Callithrix jacchus*) under laboratory conditions. *Abstracts, XIth Congress of the International Primatological Society*, Göttingen.
- Rylands, A. B. 1989. Sympatric Brazilian callitrichids: the black-tufted-ear marmoset, *Callithrix kuhli*, and the golden-headed lion tamarin, *Leontopithecus chrysomelas*. *J. Hum. Evol.* 18: 679-695.
- Scanlon, C. E., Chalmers, N. R. and Monteiro da Cruz, M. A. O. 1988. Changes in the size, composition and reproductive condition of wild marmoset groups (*Callithrix jacchus jacchus*) in north east Brazil. *Primates* 29: 295-305.
- Schröpel, M. 1994. Zehn Jahre Krallenaffenhaltung im Zoologischen Garten Magdeburg. *Felis* 12: 44-54.
- Soini, P. 1982. Primate conservation in Peruvian Amazonia. *Int. Zoo Yb.* 22: 37-47.
- Soini, P. 1988. The pygmy marmoset, genus *Cebuella*. In: *Ecology and Behavior of Neotropical Primates*. Vol. 2, R. A. Mittermeier, A. B. Rylands, A. F. Coimbra-Filho and G. A. B. da Fonseca (eds.), pp.79-129. World Wildlife Fund, Washington, D. C.
- Soini, P. 1993. The ecology of the pygmy marmoset, *Cebuella pygmaea*: Some comparisons with two sympatric tamarins. In: *Marmosets and Tamarins: Systematics, Behaviour, and Ecology*, A. B. Rylands (ed.), pp.257-261. Oxford University Press, Oxford.
- Stevenson, M. F. and Rylands, A. B. 1988. The marmosets, genus *Callithrix*. In: *Ecology and Behavior of Neotropical Primates*, Vol. 2., R. A. Mittermeier, A. B. Rylands, A. F. Coimbra-Filho and G. A. B. da Fonseca (eds.), pp.131-222. World Wildlife Fund, Washington, D. C.
- Tardif, S. D. 1984. Social influences on sexual maturation of female *Saguinus oedipus*. *Am. J. Primatol.* 6: 199-209.
- Ziegler, T. E., Scholl, S. A., Scheffler, G., Haggerty, M. A. and Lasley, B. L. 1989. Excretion of oestrone, oestradiol and progesterone in the urine and feces in the female cotton-top tamarin (*Saguinus oedipus oedipus*). *Am. J. Primatol.* 17: 185-195.

THE EFFECT OF RAINFALL SEASONALITY ON THE GEOGRAPHIC DISTRIBUTION OF NEOTROPICAL PRIMATES

R. Pastor-Nieto
D. K. Williamson

Introduction

Studies carried out on the geographic distribution and ecology of New World Primates (for example, Pagel *et al.*, 1991; Ross, 1992; Rosenberger, 1992; Strier, 1992; Ford, 1994) all find that environmental variables such as temperature and rainfall have an important effect on the morphological adaptations and behavioral ecology of the platyrrhine primates.

There is a general relationship between overall body size and ecological, behavioral and physiological traits in mammals (for example, Calder, 1984; Clutton-Brock and Harvey, 1983; Eisenberg, 1981; McNab, 1987; Peters, 1983; Schmidt-Nielsen, 1984). In primates there is a consistent relationship between body size and feeding ecology (Ford and Davis, 1992). Body size by itself, therefore, is a useful predictor of a species' adaptations (Damuth and McFadden, 1990), and appears to be related to numerous life-history variables (Clutton-Brock and Harvey, 1983; Harvey and Clutton-Brock, 1985).

Platyrrhines are thought ideal for examining morphological adaptations to diet, because of their monophyletic origin, universal arboreality, and variation in food prefer-

Table 1. Mean body weights (adapted from Ford and Davis, 1992) of the platyrrhine primates, and their presence in the phytogeographic regions (Rylands *et al.*, 1995). *CCCH = Cerrado/Caatinga/Chaco; AM = Amazonia; SA = Southern Andes; MA = Middle America; NVC = Northern Venezuela-Colombia; NA = Northern Andes; AF = Atlantic forest.

Species	Male weight (g)	Female weight (g)	Mean weight (g)	Phytogeographic region
<i>Alouatta belzebul</i>	7270.0	5525.0	6397.5	AM
<i>Alouatta caraya</i>	6800.0	4605.0	5702.5	CCCH
<i>Alouatta fusca</i>	6175.0	4550.0	5362.5	AF
<i>Alouatta palliata</i>	7150.0	5350.0	6250.0	MA, NA
<i>Alouatta pigra</i>	11352.0	6434.0	8893.0	MA
<i>Alouatta seniculus</i>	7200.0	5600.0	6400.0	SA, MA, NVC
<i>Aotus azarae</i>	933.0	953.0	943.0	CCCH, AM
<i>Aotus lemurinus</i>	955.0	968.0	961.5	MA, NA
<i>Aotus nancymae</i>	923.0	940.0	931.5	AM
<i>Aotus trivirgatus</i>	920.0	950.0	935.0	AM
<i>Ateles belzebuth</i>	6200.0	5800.0	6000.0	AM, NVC, NA
<i>Ateles fusciceps</i>	8890.0	8800.0	8845.0	MA, NVC, NA
<i>Ateles geoffroyi</i>	8210.0	7456.0	7833.0	MA
<i>Ateles paniscus</i>	7460.0	9750.0	8605.0	AM
<i>Brachyteles arachnoides</i>	12125.0	9450.0	10787.5	AF
<i>Cacajao calvus</i>	3450.0	2880.0	3165.0	AM
<i>Cacajao rubicundus</i>	3450.0	2880.0	3165.0	AM
<i>Cacajao melanocephalus</i>	3450.0	2880.0	3165.0	AM
<i>Callicebus brunneus</i>	854.0	805.0	829.5	AM
<i>Callicebus caligatus</i>	1098.0	1075.0	1086.5	AM
<i>Callicebus cinerascens</i>	1098.0	1075.0	1086.5	AM
<i>Callicebus cupreus</i>	1012.0	1119.0	1065.5	AM
<i>Callicebus donacophilus</i>	1098.0	1075.0	1086.5	CCCH, AM
<i>Callicebus dubius</i>	1098.0	1075.0	1086.5	AM
<i>Callicebus hoffmannsi</i>	1098.0	1075.0	1086.5	AM
<i>Callicebus moloch</i>	1000.0	860.0	930.0	AM
<i>Callicebus oenanthe</i>	1098.0	1075.0	1086.5	SA
<i>Callicebus olallae</i>	1098.0	1075.0	1086.5	AM
<i>Callicebus personatus</i>	1325.0	1285.0	1305.0	AF
<i>Callicebus torquatus</i>	1300.0	1307.0	1303.5	AM
<i>Callimico goeldii</i>	640.0	530.0	585.0	AM
<i>Callithrix argentata</i>	357.0	320.0	338.5	AM
<i>Callithrix geoffroyi</i>	290.0	190.0	240.0	AF
<i>Callithrix humeralifera</i>	280.0	310.0	295.0	AM
<i>Callithrix jacchus</i>	256.0	236.0	246.0	CCCH, AF
<i>Cebuella pygmaea</i>	130.0	126.0	128.0	AM
<i>Cebus albifrons</i>	2480.0	1814.0	2147.0	AM, SA, NVC, NA
<i>Cebus apella</i>	3050.0	2385.0	2717.5	CCCH, AM, SA, NVC, AF
<i>Cebus capucinus</i>	3868.0	2666.0	3267.0	MA, NVC, NA
<i>Cebus olivaceus</i>	2974.0	2395.0	2684.5	AM, NVC
<i>Chiropotes albinasus</i>	3020.0	2510.0	2765.0	AM
<i>Chiropotes satanas</i>	3100.0	2600.0	2850.0	AM
<i>Lagothrix flavicauda</i>	8335.0	5750.0	7042.5	SA
<i>Lagothrix lagothricha</i>	8335.0	5750.0	7042.5	AM
<i>Leontopithecus chrysomelas</i>	620.0	535.0	577.5	AF
<i>Leontopithecus chrysopygus</i>	614.0	557.0	585.5	AF
<i>Leontopithecus rosalia</i>	607.0	578.0	592.5	AF
<i>Pithecia irrorata</i>	2010.0	1875.0	1942.5	AM
<i>Pithecia monachus</i>	2795.0	1900.0	2347.5	AM
<i>Pithecia pithecia</i>	1732.0	1515.0	1623.5	AM
<i>Saguinus bicolor</i>	430.0	430.0	430.0	AM
<i>Saguinus fuscicollis</i>	387.0	403.0	395.0	AM
<i>Saguinus imperator</i>	400.0	400.0	400.0	AM
<i>Saguinus inustus</i>	423.0	454.0	438.5	AM
<i>Saguinus labiatus</i>	451.0	465.0	458.0	AM
<i>Saguinus leucopus</i>	490.0	490.0	490.0	AM
<i>Saguinus midas</i>	586.0	432.0	509.0	AM
<i>Saguinus mystax</i>	577.0	560.0	568.5	AM
<i>Saguinus nigricollis</i>	470.0	480.0	475.0	AM
<i>Saguinus oedipus</i>	411.0	430.0	420.5	AM
<i>Saguinus tripartitus</i>	423.0	454.0	438.5	AM
<i>Saimiri boliviensis</i>	1015.0	700.0	857.5	CCCH, AM
<i>Saimiri oerstedii</i>	829.0	695.0	762.0	MA
<i>Saimiri sciureus</i>	852.0	675.0	763.5	AM, NVC
<i>Saimiri ustus</i>	910.0	795.0	852.5	AM
<i>Saimiri vanzolinii</i>	950.0	650.0	800.0	AM




Figure 1. The geographic ranges of a large (*Alouatta*) and a small (*Cebuella*) platyrrhine, overlaid with the seven phytogeographic regions of Gentry (1982).

ences (Anapol and Lee, 1994). Differentiation of body size is a significant factor in the partitioning of platyrrhine diets and foraging strategies (for example, Tamarin *et al.*, 1984; Janson and Boinski, 1992; Garber, 1992; Rosenberger and Strier, 1989; Strier, 1992).

There are currently three main hypotheses, based on ecological determinants, to explain the maximum body-weight thresholds of platyrrhines.

Terborgh and van Schaik (1987) suggested that the fruit and leafing cycles in the Neotropics are 'in-phase', therefore the species have to adapt to a seasonal environment. This leaves two options; either to specialize as folivores, or to exploit insects in periods of fruit scarcity.

Fleming *et al.* (1987) suggested that the low diversity of fleshy fruits in the Neotropics, in combination with a greater variety of fruit species, would have favored the evolution of small-bodied frugivores.

The 'brittle-branch' hypothesis (Cristoffer, 1987) proposed that the Neotropical forests had a more 'fragile' vegetation structure than its Paleotropical counterpart (Emmons and Gentry, 1983) and would have limited the evolution of large-bodied arboreal vertebrates.

All of these theories converge on one simple statement: the body size of Neotropical primates seems to be indirectly affected by climate. However, such theories do not explain the current geographical distribution of species. In this context we make two assumptions:

The larger platyrrhines, such as the atelines are distributed more evenly in the Neotropics because they are buffered against long periods of reduced food availability during the dry season (Boyce, 1979; Lindstedt and Boyce, 1985).

Table 2. Climatic variables calculated from long term records from randomly sampled weather stations (Wernstedt, 1972) for each of the phyto-geographic regions (adapted from Gentry, 1982). CCCH = Cerrado/Caatinga/Chaco; AM = Amazonia; SA = Southern Andes; MA = Middle America; NVC = Northern Venezuela-Colombia; NA = Northern Andes; AF = Atlantic forest.

Phytogeographic region	Climatic variables*						
	n	T	P	RDV	Z	MO<50	P>2T
CCCH	38	21.45	1116.62	0.74	0.89	3.65	8.68
AM	30	25.08	2118.01	0.71	0.90	2.18	9.79
SA	8	23.46	1407.29	0.81	0.89	3.13	10.67
MA	20	25.73	2876.04	0.82	0.89	2.00	9.90
NVC	11	22.60	2198.11	0.79	0.90	2.55	9.36
NA	7	21.31	772.75	0.91	0.85	6.00	6.57
AF	7	21.21	1439.75	0.79	0.90	2.71	9.57
CCCH/AF	**	21.33	1302.18	0.78	0.90	3.18	9.12
CCCH/AM	**	23.26	1641.31	0.74	0.89	2.19	9.24
AF/AM	**	23.14	1778.88	0.75	0.90	2.45	9.68
AM/SA	**	23.18	1443.37	0.81	0.87	4.09	8.18
AM/NA	**	20.34	1444.83	0.82	0.89	3.71	9.90
AM/NVC	**	23.83	2158.08	0.73	0.90	2.36	9.58
NA/MA	**	23.51	1854.58	0.87	0.87	4.00	8.24
NA/NVC	**	21.95	1485.42	0.87	0.87	4.27	7.97
MA/NVC	**	234.16	2357.07	0.89	0.89	2.27	9.63

*n = number of weather stations sampled; T = mean annual temperature; P = mean annual precipitation (°C); RDV = rainfall diversity index; Z = Simpson's index of diversity; MO<50 = number of months where mean rainfall is greater than 50mm; P>2T = number of months where rainfall (mm) is greater than two times the temperature (°C).

**No sample size given since means for the overlapping sub-regions were calculated from the means of two neighboring regions.

Smaller species, such as the callitrichids, are more restricted in their geographic distribution due to their nutritional requirements (see Fleagle, 1988).

To illustrate this observation two genera were selected: a larger ateline with a wide distribution (*Alouatta* spp.), and the smallest platyrrhine genus (*Cebuella* spp.) which has a relatively restricted distribution (see Fig 1).

In this paper we test whether rainfall seasonality has an effect on the geographic distribution of platyrrhine species.

Methods

Southern and Middle America can be divided into nine phytogeographic regions (Gentry 1982), which give an accurate description of Neotropical vegetation diversity. In a review of platyrrhine distribution, Rylands *et al.* (1995) listed the species occurring in seven phytogeographic regions, adapted from Gentry (1982) (see Fig. 1). They did not consider two regions, the Caribbean, because no primates occur there, and the Guyana sub-region, which is subsumed into the Amazonia region. We use the species list of Rylands *et al.* (1995) because of the detailed information on distributions provided (see Table 1).

Indexing habitat productivity and rainfall seasonality

The main climate variables influencing plant and animal life are water availability and temperature, and can be used alone to characterize vegetation patterns and plant productivity globally (Walter, 1979; Box, 1981; Le Houerou

Table 3. The smallest species for each weight category, rainfall diversity (RDV) and mean annual temperature for each of Gentry's (1982) phylogeographic regions. P values and r^2 values listed for weight category versus RDV regressions.

Phylogeographic Regions	Body weight categories (g)						
	< 500	600-1000	1000-1500	1500-2500	2500-3300	3300-6400	6400-11000
AM	128.0	807.0	1004.0	1732.0	3165.0	6200.0	7621.0
AF	246.0	1933.0	1004.0	*	2675.0	5362.0	10788.0
CCCH	246.0	607.0	1004.0	*	2171.5	5702.0	*
MA	420.5	955.0	*	*	3267.0	6250.0	7833.0
NVC	420.5	955.0	1065.5	*	2675.0	6400.0	8605.0
NA	490.0	955.0	1303.5	*	2675.0	*	7621.0
SA	*	955.0	*	*	2675.0	6400.0	7042.5
Diet***	I, Fr, G	I, Fr, Pr, ScFr	ScFr	*	ScFr	Fol	Fr, Fol
p values	0.001	0.01	0.04	**	0.5	0.28	0.81
r^2	0.87	0.62	0.93	**	0.28	0.20	0.06

* No species of that weight category for that particular phylogeographic region.

** No statistical tests conducted.

*** I = insectivore, Fr = frugivore, G = gummivore, ScFr = sclerocarpic-frugivore, Fol = folivore, Pr = vertebrate predator.

and Popov, 1981). However, plant productivity can be indexed indirectly in a variety of ways; plant evapotranspiration being the most accurate. Potential evapotranspiration (PET) measures the potential amount of water released into the atmosphere through plant evaporation, respiration and transpiration. In the absence of data to compute evapotranspiration, it has been suggested (Bagnouls and Guassen, 1953; Walter and Leith, 1967) that the number of months where precipitation (P), in millimeters, is less than two times the mean annual temperature (t), in degrees Celsius ($P > 2t$) is an excellent substitute for PET. This is shown by the high correlation between the two measures ($r^2 = 0.938$, $p = 0.00$; Williamson, 1997). The advantage of the $P > 2t$ index is that the data are readily available.

Seasonality of Gentry's phylogeographic regions

Gentry (1982) characterized the phylogeographic regions in terms of plant species richness, (canopy trees and lianas; epiphytes and palmettos). Clinebell *et al.* (1995) further suggested that Neotropical tree and liana species' richness are best explained by annual rainfall and rainfall seasonality. In this analysis we characterized each of the phylogeographic regions in terms of climate, with a view

to predicting habitat productivity. There is also strong evidence that insect abundance, and fruiting and leafing phenology is highly correlated with rainfall seasonality (Wolda, 1978; Terborgh, 1983; Poulin *et al.*, 1992).

Weather stations from a worldwide climatic data source (Wernstedt, 1972) were chosen for each of the phylogeographic regions. They were selected focusing on the sampling areas in each region around known field study sites where the primates are known to live. Mean values for the climatic variables were then calculated for each of the phylogeographic regions.

Many of the primates occur in more than one phylogeographic region. These regions of overlap were considered as discrete sub-regions. Those species in the overlapping sub-regions were considered to be at the limits of their distribution, and therefore assumed to be in the most ecologically constrained part of their range. In support of this approach, it has been suggested (Strier, 1992), that one way to further distinguish between the effects of seasonality on dietary strategies (and by inference body size), is to examine data on sympatric taxa. The same climatic variables were recorded for the regions of overlap.

The following climatic variables were recorded from each of the weather stations; mean annual temperature ($^{\circ}\text{C}$); mean annual rainfall (mm), number of months with less than 50 mm ($mo < 50$); the number of months where mean monthly rainfall (mm) was greater than twice the mean monthly temperature ($^{\circ}\text{C}$); and ($P > 2t$). Two indices of rainfall diversity were calculated. The first index was Z (Simpson's index of rainfall diversity). Simpson's is an index of the proportional distribution of total rainfall across the 12 months of the year, and varies from $Z = 0$ (completely uneven) to $Z = 1$ (completely even). Simpson's index of diversity (Peet, 1974), which is derived from the species diversity literature (Magurran, 1988), has been questioned (Bronikowski and Webb, 1996; but see Williamson and Dunbar, in press), since it does not take dry months into account. There is as yet no single climatic index applicable across all habitat types. Some indices provide insufficient variance between values of the




Figure 2. The smallest platyrrhine species' body weight in each of the phylogeographic regions (Gentry, 1982), regressed against rainfall diversity (RDV). CCCH = Cerrado/Caatinga/Chaco; AM = Amazonia; SA = Southern Andes; MA = Middle America; NVC = Northern Venezuela/Colombia; NA = Northern Andes; AF = Atlantic forest.

seasonality index for statistical tests; for example, when rainfall is very low or even. We used, therefore, the index of the temporal distribution (RDV) of rainfall (Williamson, 1997). RDV is calculated by finding the difference between all possible pair-wise comparisons of the 12 mean monthly rainfall values, then finding the mean of these differences. This statistic is based on the exact randomization test (Sokal and Rohlf, 1981; p.788). RDV records the opposite of Z, a high value indicates a more seasonal environment, a low value a less seasonal environment. The climate data for the regions and overlapping regions is shown in Table 2.

Body weight categories

We divided the platyrrhines into body weight categories that would reflect finer-grained dietary adaptations, since body size and dietary quality are related (Fleagle, 1988; Anapol and Lee, 1994). To test whether rainfall seasonality has an effect on the geographic distribution of platyrrhines, the smallest species was chosen for each of the seven body weight categories, for each of Gentry's regions (Table 3). The smallest species were selected since they would be expected to be more constrained in their distribution, and because small species they are less able to buffer themselves against unpredictable food availability in fluctuating environments.

Results

Initial diagnostic Spearman rank correlations showed RDV to be the only climatic variable significantly correlated with body weight, so only those results will be presented here.

Table 4. Genera inhabiting each phytogeographic region, and those occupying regions overlapping phytogeographic regions. MA = Middle America; NA = Northern Andes; NVC = Northern Venezuela-Colombia; SA = Southern Andes; AM = Amazonia; AF = Atlantic Forest; CCCH = Cerrado/Caatinga/Chaco.

Phytogeographic region	Genera
MA	<i>Saguinus, Aotus, Saimiri, Cebus, Alouatta, Ateles</i>
NA	<i>Saguinus, Aotus, Callicebus, Cebus, Alouatta, Ateles, Lagothrix</i>
NVC	<i>Saguinus, Aotus, Callicebus, Saimiri, Cebus, Alouatta, Lagothrix</i>
SA	<i>Aotus, Callicebus, Cebus, Alouatta, Lagothrix</i>
AM	<i>Cebuella, Callithrix, Saguinus, Aotus, Callicebus, Saimiri, Cebus, Pithecia, Chiropotes, Cacajao, Alouatta, Ateles, Lagothrix</i>
AF	<i>Callithrix, Leontopithecus, Callicebus, Cebus, Alouatta, Brachyteles</i>
CCCH	<i>Callithrix, Aotus, Callicebus, Cebus, Alouatta</i>
Overlapping regions	
CCCH/AF	<i>Callithrix, Callicebus, Cebus, Alouatta</i>
CCCH/AM	<i>Callithrix, Aotus, Callicebus, Cebus, Alouatta</i>
AF/AM	<i>Callithrix, Callicebus, Cebus, Alouatta</i>
AM/NA	<i>Saguinus, Aotus, Callicebus, Saimiri, Cebus, Alouatta, Ateles</i>
AM/SA	<i>Aotus, Callicebus, Cebus, Alouatta, Lagothrix</i>
AM/NVC	<i>Saguinus, Aotus, Saimiri, Cebus, Alouatta, Ateles</i>
NA/MA	<i>Saguinus, Aotus, Cebus, Alouatta</i>
NA/NVC	<i>Saguinus, Aotus, Callicebus, Cebus, Alouatta, Ateles</i>
NA/NVC	<i>Saguinus, Aotus, Saimiri, Cebus, Alouatta, Ateles</i>

Body weight relationships with climate

The body weight of the smallest species in each of the seven phytogeographic regions was positively correlated with RDV ($r^2 = 0.70$, $F = 2.19$, $p = 0.01$), (Fig. 2). The regions containing the smallest species had the most even spread of rainfall (low RDV, or non-seasonal environments).

Body weight categories were regressed against RDV (Table 3, Fig. 3). As body weight increases, the significance level of the regression slope decreases (Table 3), being non-significant from the 2300 g category and upwards (Figs. 3: D, E. and F.). The most highly significant regression is that for the category <500g ($r^2 = 0.87$, $p = 0.001$). This corresponds to "Kay's threshold" (Kay, 1975) which defines the boundary between insectivorous (<500g) and folivorous primates (>500g).

Discussion

This study suggests that rainfall seasonality acts as a selective force on the distribution of body weights in Neotropical primates. The results of this study summarize what has already been suggested by other authors (for example, Cristoffer, 1987; Ross, 1992; Kinzey, 1994), that the adaptive morphological changes in body size are affected by environmental factors. In this paper we have analyzed in detail how environmental factors characterize rainfall seasonality and hence habitat productivity in each of Gentry's phytogeographic regions.

The most important result presented in this paper is the very significant effect that rainfall seasonality has on the geographic boundaries of the smallest species and how, in very seasonal habitats, only larger platyrrhines are found. Species' with body weights of less than 500g can only be distributed in the Amazonian region, which has a tendency to be less seasonal, favoring year-round food availability.

The smallest body weight category plotted against rainfall seasonality (RDV) (Fig. 3: A; Table 3), has the most highly significant regression slope of all the body weight categories ($r^2 = 0.87$, $p = 0.001$). This result corresponds to "Kay's threshold" (Kay 1975, 1984), and applies to the entire Order Primates. Smaller animals require a higher protein content in their diet, and therefore have a higher

Table 5. Mean body weights for each of the platyrrhine genera.

Genus	Mean weight (g)
<i>Alouatta</i>	6500.92
<i>Aotus</i>	942.75
<i>Ateles</i>	7820.75
<i>Brachyteles</i>	10787.5
<i>Cacajao</i>	3165
<i>Callicebus</i>	1086.58
<i>Callimico</i>	585
<i>Callithrix</i>	279.88
<i>Cebuella</i>	128
<i>Cebus</i>	2704
<i>Chiropotes</i>	2807.5
<i>Lagothrix</i>	7042.5
<i>Leontopithecus</i>	585.17

basal metabolic rate (BMR), and consequently higher energetic needs (reviewed in Ford and Davis, 1992). Kay (1984) stated that no primarily insectivorous primates (>30-40% of their diet composed of insects) would weigh more than 700g. Significantly, this body weight threshold also correlates with twinning and monogamy (Dunbar, 1995a, 1995b), features only found in callitrichids. On the other hand, for the middle-sized and larger platyrrhines (>1 kg), troop fragmentation appears to be a facultative adjustment to food scarcity (Kinzey and Cunningham, 1984).

Rosenberger (1992) suggested that small and large platyrrhines have overcome the problem of seasonality in two different ways:

1. Smaller species, which require a high energy diet, will shift to alternative food resources (e.g., gums or insects)

in the dry season when fruit is scarce. For example, *Saguinus fuscicollis* switches from fruit to predominantly nectar in the dry season (Terborgh and Stern, 1987), and thus avoids competition with larger primates by feeding on specialized resources (Garber, 1992). Alternatively Peres (1989) suggests, that small species switch to different feeding locations, where insects are more likely to be concentrated in the dry season, such as gumlicks and knot-holes. We need to be cautious, therefore, in suggesting that resources for the smaller primates are fewer in the dry season.

2. Larger platyrrhine primates have solved the problem of seasonality with folivory. Leaves are available year-round, although they are a low-quality food resource that requires special adaptations to overcome the problems of its digestion (Rosenberger, 1992). Large body size, an adaptation




Figure 3. The smallest platyrrhine primate in each of the body weight categories (see Table 3) in each phytogeographic region against rainfall diversity (RDV). A. less than 500 g (dotted line indicates Kay's threshold); B. 600-1000 g; C. 1000-1500 g; D. 2500-3300 g; E. 3300-6400 g; F. 6400-11000 g.

to folivory, buffers against harsh seasonal habitats. This is supported by recent discoveries of giant extinct pitheciines (Hartwig, 1995; Hartwig and Cartelle, 1996), weighing as much as 25 kg. Their extreme large size may be an adaptation to competition for resources in a harsh habitat, brought on by climate changes in the Pleistocene.

Low rainfall seasonality habitats evidently permit the survival of a wider variety of primate species, but there is a need to understand better the environmental factors that underlie the biogeographic distribution of species if we are to devise effective conservation strategies.

Acknowledgments

We are grateful to Prof. R. Dunbar for his comments on earlier drafts, to D. Tzivilis for assistance with scanning, and to C. Lowen and C. Underwood for discussions on the work in progress. R. Pastor-Nieto has a grant from The National Council of Science and Technology of Mexico (CONACYT). D. K. Williamson is grateful for support from the Natural Environment Research Council (NERC) and the Leverhulme Trust.

R. Pastor-Nieto, Coordinador Proyecto Hampolol, Universidad Autonoma de Campeche, ECOMAT, Avenida Universidad s/n, Campeche, Campeche, México, and **D. K. Williamson**, Hominid Palaeontology Research Group, Department of Human Anatomy and Cell Biology, University of Liverpool, New Medical School, Ashton Street, Liverpool L69 3GE, UK.

References

- Anapol, F. and Lee, S. 1994. Morphological adaptation to diet in platyrrhine primates. *Am. J. Phys. Anthropol.* 94: 239-261.
- Boyce, M. S. 1979. Seasonality and patterns of natural selection for life histories. *Am. Nat.* 114, 569-583.
- Box, E. O. 1981. *Macroclimate and Plant Forms: An Introduction to Predictive Modeling in Phytogeography*. W. Junk, The Hague.
- Bronikowski, A. M. and Webb, C. 1996. Appendix: A critical examination of rainfall variability measures used in behavioral ecology studies. *Behav. Ecol. Sociobiol.* 39: 27-30.
- Calder, W. A. 1984. *Size, Function and Life History*. Harvard University Press, Cambridge, MA.
- Clinebell II, R.R., Philips, O. L., Gentry, A. H., Stark, N. and Zuuring, H. 1995. Prediction of neotropical tree and liana species richness from soil and climatic data. *Biodiversity and Conservation* 4: 56-90.
- Clutton-Brock, T. H. and Harvey, P. 1983. The functional significance of body size among mammals. In: *Advances in the Study of Mammalian Behavior*, J. F. Eisenberg and D. G. Kleiman (eds.), pp.632-663. Special Publication of the American Society of Mammalogists, No. 7. Academic Press, New York.
- Coe, M. J., Cumming, D. H. and Philipson, J. 1976. Biomass and production of large herbivores in relation to rainfall and primary production. *Oecologia* 22: 342-354.
- Cristoffer, C. 1987. Body size differences between New World and Old World, arboreal, tropical vertebrates: cause and consequences. *J. Biogeog.* 14: 154-172.
- Damuth, J. and McFadden, B. J. 1990. *Body Size in Mammalian Paleobiology: Estimation and Biological Implications*. Cambridge University Press, Cambridge.
- Dickinson, R. E. (ed.). 1987. *The Geophysics of Amazonia: Vegetation and Climate Interactions*. John Wiley and Sons, New York.
- Dunbar, R. I. M. 1995a. The mating system of callitrichid primates: I. Conditions for the evolution of pair bonding and twinning. *Anim. Behav.* 50: 1057-1070.
- Dunbar, R. I. M. 1995b. The mating system of callitrichid primates: II. The impact of helpers. *Anim. Behav.* 50: 1071-1089.
- Emmons, L. H. and Gentry, A. H. 1983. Tropical forest structure and the distribution of gliding and prehensile tailed vertebrates. *Am. Nat.* 124: 513-524.
- Eisenberg, J. F. 1981. *The Mammalian Radiations: An Analysis of Trends in Evolution, Adaptation and Behavior*. Chicago University Press, Chicago.
- Fleagle, J. G. 1988. *Primate Adaptation and Evolution*. Academic Press, San Diego.
- Fleming, T. H., Breitwisch, R. and Whitesides, G. H. 1987. Patterns of tropical vertebrate frugivore diversity. *Ann. Rev. Ecol. Syst.* 18: 91-109.
- Ford, S. M. 1992. Evolution of sexual dimorphism in body weight in platyrrhines. *Am. J. Primatol.* 34: 221-224.
- Ford, S. M. and Davis, L. C. 1992. Systematics and body size: Implications for feeding adaptations in New World Monkeys. *Am. J. Phys. Anthropol.* 88: 415-468.
- Garber, P. A. 1992. Vertical clinging, small body size, and the evolution of feeding adaptations in the Callitrichidae. *Am. J. Phys. Anthropol.* 88: 469-482.
- Gentry, A. H. 1982. Neotropical floristic diversity: phyto-geographical connections between Central and South America, Pleistocene climatic fluctuations, or an accident of the Andean orogeny? *Ann. Missouri Bot. Gard.* 69: 557-593.
- Hartwig, W. C. 1995. A giant New World monkey from the Pleistocene of Brazil. *J. Hum. Evol.* 28: 189-195.
- Hartwig, W. C. and Cartelle, C. 1996. A complete skeleton of the giant South-American primate *Protopithecus*. *Nature, Lond.* 381: 307-311.
- Harvey, P. H. and Clutton-Brock, T. H. 1985. Life-history variation in primates. *Evolution*, 39: 559-581.
- Janson, C. H. and Boinski, S. 1992. Morphological and behavioural adaptations for foraging in generalist primates: the case of the cebines. *Am. J. Phys. Anthropol.* 88: 483-498.
- Kay, R. F. 1975. The functional adaptations of primate molar teeth. *Am. J. Phys. Anthropol.* 43: 195-216.
- Kay, R. F. 1984. On the use of anatomical features to infer foraging behavior in extinct primates. In: *Adaptations for Foraging in Nonhuman Primates*, P. S. Rodman and J. G. H. Cant (eds.), pp.21-53. Colombia University Press, New York.
- Kinzey, W. G., and Cunningham, E. P. 1994. Variability in platyrrhine social organization. *Am. J. Primatol.* 34:

- 185-198.
- Le Houerou, H. N. and Popov, G. F. 1981. An ecoclimatic classification of intertropical Africa. *Plant. Prod. Protec. Pap.* (31). F.A.O., Rome.
- Lindstedt, S. and Boyce, M. 1985. Seasonality, fasting endurance, and body size in mammals. *Am. Nat.* 125: 873-878.
- MacArthur, R. H. and Wilson, E. O. 1967. *The Theory of Island Biogeography*. Princeton University Press, Princeton, NJ.
- Magurran, A. E. 1988. *Ecological Diversity and its Measurement*. Princeton University Press, Princeton, NJ.
- MacNab, B. 1990. The physiological significance of body size. In: *Body Size in Mammalian Paleobiology: Estimation and Biological Implications*, J. Damuth and B. J. MacFadden (eds.), pp.11-23. Cambridge University Press, Cambridge.
- Murphy, P. G. 1975. Net primary productivity in tropical terrestrial ecosystems. In: *Primary Productivity of the Biosphere*, H. Leith and R. H. Whittaker (eds.), pp.217-231. Springer Verlag, Berlin.
- Pagel, M. D., May, R. D. and Collie, A. R. 1991. Ecological aspects of the geographical distribution and diversity of mammalian species. *Am. Nat.* 137: 791-815.
- Peet, R. K. 1974. The measurement of diversity. *Ann. Rev. Ecol. Syst.* 5: 285-307.
- Peres, C. A. 1989. Exudate-eating by wild golden lion tamarins, *Leontopithecus rosalia*. *Biotropica*, 21: 287-288.
- Peters, R. H. 1983. *The Ecological Implications of Body Size*. Cambridge University Press, Cambridge.
- Poulin, B., Lefebvre, G. and MacNeil, R. 1992. Tropical avian phenology in relation to abundance and exploitation of food resources. *Ecology* 73: 2295-2309.
- Reed, K. E. and Fleagle, J. G. 1995. Geographic and climatic control of primate diversity. *Proc. Natl. Acad. Sci. USA*. 92: 7874-7876.
- Ross, C. 1992. Environmental correlates of the intrinsic rate of natural increase in primates. *Oecologia* 90: 383-390.
- Rosenberger, A. L. 1992. Evolution of feeding niches in New World monkeys. *Am. J. Phys. Anthropol.* 88: 525-562.
- Rosenberger, A. L. and Strier, K. B. 1992. Adaptive radiation of ateline primates. *J. Hum. Evol.* 18: 717-750.
- Rylands, A. B., Mittermeier, R. A. and Rodríguez-Luna, E. 1995. A species list for the New World primates (Platyrrhini): Distribution by country, endemism, and conservation status according to the Mace-Lande system. *Neotropical Primates* 3 (suppl.): 113-164.
- Schmidt-Nielsen, K. 1990. *Animal Physiology: Adaptation and Environment*. 4th ed. Cambridge University Press, Cambridge.
- Sokal, R. R. and Rohlf, F. J. 1981. *Biometry: The Principles and Practice of Statistics in Biological Research*, 2nd ed. W. H. Freeman, New York.
- Strier, K. B. 1992. Aeline adaptations: behavioral strategies and ecological constraints. *Am. J. Phys. Anthropol.* 88: 515-524.
- Temerin, L. A., Wheatley, B. P. and Rodman, P. S. 1984. Body size and foraging in primates. In: *Adaptations for Foraging in Non-Human Primates*, P. S. Rodman and J. G. H. Cant (ed.), pp.215-248. Columbia University Press, New York.
- Terborgh, J. 1983. *Five New World Primates: A Study in Comparative Ecology*. Princeton University Press, Princeton, NJ.
- Terborgh, J. and Stern, M. 1987. The surreptitious life of the saddle-backed tamarin. *Am. Sci.* 75: 260-269.
- Walter, H. and Leith, H. 1967. *Climate Diagram World Atlas*. Gustav Fischer, Jena.
- Wernstedt, F. L. 1972. *World Climatic Data*. Climatic Data Press, Lemont.
- Williamson, D. K. 1997. Primate Socioecology: Development of a Conceptual Model for the Early Hominids. Unpublished Ph.D. thesis, University College London, London.
- Williamson, D. K. and Dunbar R. I. M. In press. Energetics, time budgets and group size. In: *Comparative Primate Socioecology*, P. C. Lee (ed.). Cambridge University Press, Cambridge.
- Wolda, H. 1978. Seasonal fluctuation in rainfall, food and abundance of tropical insects. *J. Anim. Ecol.* 47: 369-381.
- Wolfheim, J. 1983. *Primates of the World*. University of Washington Press, Seattle.

THE SQUIRREL MONKEY BREEDING COLONY OF THE PASTEUR INSTITUTE, CAYENNE, FRENCH GUIANA

Benoît de Thoisy
Hugues Contamin

Introduction

Squirrel monkeys, especially *Saimiri sciureus* and *S. boliviensis*, are animal models widely used in biomedical research; mainly in pharmacology, toxicology, cancerology, nutrition, cardiovascular diseases and neurology (Mittermeier *et al.*, 1994). Numerous captive breeding colonies have been established in different parts of the world. However, reproduction of captive squirrel monkeys is often inconsistent, giving rise to disappointing breeding results. The aim of this article is to briefly present the 15-year-old colony of the Pasteur Institute of French Guiana, emphasizing efforts to ensure the well-being of the monkeys and, being a biomedical research institution, the results obtained using this non-human primate model.

In 1978, The Pasteur Institute of French Guiana, belonging to a network of international Pasteur Institutes, decided to initiate a long-term research program on human malaria. Although not the natural host of either *Plasmodium vivax* or *P. falciparum*, *Saimiri sciureus* was chosen as a model because it is sensitive to experimental infection (Gysin, 1991). Moreover, squirrel monkeys naturally